arXiv:gr-qc/0109060v2 28 Sep 2001

X-matter induced cosmological scenarios in the relativistic theory of gravity

V. L. Kalashnikov

7th February 2008

Institut für Photonik, TU Wien, Gusshausstr. 27/387, A-1040 Vienna,
Austria, tel:+43-1-58801-38723, fax:+43-1-58801-38799,
e-mail:vladimir.kalashnikov@tuwien.ac.at

Abstract

It was shown, that the presence of the so-called X-matter with an equation of state, which lies between limits of the strong and weak energy conditions, allows the variety of the cosmological scenarios in the relativistic theory of gravity. In spite of the fixed negative sign of the cosmological term in the field equations with massive graviton, it is possible to obtain the solutions with accelerated and complicated loitering expansion of the universe. The numerical estimation of the universe's age agrees with the modern observational data if the upper limit of the graviton's mass is 10^{-71} g

1 Introduction

The so-called relativistic theory of gravity (RTG), which was inspired by A.A. Logunov with co-authors [1, 2], is the alternative to Einstein's general theory of relativity (GR) and differs from it in the cardinal point: the gravitation is considered as the tensor field of the Faraday-Maxwell's type in the flat Minkowski spacetime. The existence of the flat reference background metric in RTG allows of the unambiguity and clarity of the obtained solutions, which are similar to ones in GR but to have the property of the trivial extendibility and topological simplicity. The lasts exclude the existence of the singularity, black holes, wormholes, topological defects etc. The structure of the theory is tougher due to existence of the additional field equation on background metric, which is analog of the harmonic coordinate equation in GR and is introduced in order to exclude the spins 1 and 0 for gravitons. The additional advantages of RTG are the existence of the conservation lows and the possibility of the canonical quantization.

Table 1: Cosmological parameters [5]

cosmological parameters	observational data
H_0	68±6
Ω_K	1.1±0.12
Ω_M	0.35±0.1
Ω_X	0.65±0.15
τ_0	12.7±3
w	≤ -0.6

All experimental consequences of RTG are the same as ones in GR for experiments in solar system. But derived in [3, 4] cosmological scenario differs essentially: RTG permits only flat global spacetime universe, critical density of the matter and recollapse of universe in the case of nonzero mass of graviton. The predictions of the universe's flatness as well as the absence of the cosmological singularity in the case of nonzero mass of graviton are the advantages of the theory. But the decelerated character of the cosmological expansion and critical density of matter don't agree with the modern observational data.

The modern estimations of the basic cosmological parameters are presented in the Table 1 (H_0 is the Hubble constant in $km\ s^{-1}\ Mps^{-1}$, $\Omega_{K,M,X}$ are the normalized density parameters for the curvature, matter and cosmological constant, respectively, τ_0 is the minimal estimation of the universe's age corresponding to age of oldest globular clusters in Gyr, w is the parameter of the equation of state for the dominant energy (see below)).

The domination of Ω_X provides the accelerated expansion of the universe because of the acceleration parameter $\frac{\ddot{a}}{aH^2}\Big|_0 = \Omega_X - \frac{\Omega_M}{2} > 0$ (a is the scaling factor, H is the Hubble function, dote denotes the derivative on the time, the values are considered at the present moment) [6, 7].

In this work we demonstrate the possibility of variety of cosmological scenarios in RTG with the so-called "dark energy" (X-matter) term, that adjusts the RTG predictions with the modern astronomical data.

2 Model

We can not introduce the cosmological constant Λ with a positive sign in RTG aimed to describe the "repulsive" action of the vacuum without modification of the field equations. Indeed the field equations containing Λ and terms describing the nonzero mass of graviton m have to produce a zero curvature of the efficient Riemannian spacetime, when the tensor of the energy-momentum of matter $T_n^m = 0$ [2]. That results in $\Lambda = -m^2$ and, as consequence, "attracting" action of the vacuum (or zero cosmological constant for the massless graviton). As it was shown in [8], there is the simple possibility to modification of the basic field equations due to introduction of the massive scalar field that results in the inflation scenario and accelerated expansion of the universe. Such approach is elegant, but here we will consider the pure phenomenological alternative taking into consideration "dark energy" (X-matter), which does not cluster on the small scales, is weakly coupled to ordinary matter and has phenomenological equation of state $p_X = w_X \rho_X$ (p is the pressure, ρ is the density), where w_X lies between 0 (dust of the "usual" matter) and -1 (true cosmological constant in GR). There exist the different candidates for such X-matter. The consideration of these models is not matter of this work (for overview see [9]) and we restrict oneself to the phenomenological equation of state with constant negative w_X.

The Logunov's field equations are:

$$G_n^m - \frac{m^2}{2}(\delta_n^m + g^{mk}\gamma_{kn} - \frac{1}{2}\delta_n^m g^{pk}\gamma_{pk}) = -8\pi T_n^m, \tag{1}$$

$$D_m \tilde{g}^{mn} = 0, \tag{2}$$

where G_n^m is the Einstein's tensor defined on the efficient Riemannian spacetime with metrics g^{mn}, γ^{mn} is the metrics of the flat background Minkowski spacetime, D_m is the covariant derivative on the background spacetime, $\tilde{g}^{mn} = \sqrt{-g}g^{mn}$, $c = G = \hbar = 1$. The choice of signs in Eq. 1 is defined by the use in the calculations of the computer algebra system Maple 6 [10].

Under the assumption of homogeneity and isotropy of the efficient Riemannian spacetime (the background spacetime has a Galilean metric in our case), its interval has the following form in the spherical coordinates [2, 11]:

$$ds^2 = d\tau^2 - a(\tau)^2\left[dr^2 + r^2\left(d\theta^2 + \sin(\theta)^2 d\phi^2\right)\right], \tag{3}$$

where τ is the proper time, $a(\tau)$ is the scaling factor as function of the proper time. Hence from Eqs. (1 - 3) we have the evolutional equations:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8}{3}\pi\rho(\tau) - \frac{1}{6}\frac{m^2\left(a^2-1\right)^2\left(a^2+\frac{1}{2}\right)}{a^6}, \tag{4}$$

$$\frac{\ddot{a}}{a} = -\frac{4}{3}\pi\left(3p(\tau) + \rho(\tau)\right) - \frac{1}{6}m^2\left(1 - \frac{1}{a^6}\right). \tag{5}$$

Then, from the phenomenological equation of state $p = w\rho$ and covariant conservation low $\nabla_n T^n_m = 0$ (here ∇_n is the covariant derivative on Riemannian spacetime), we can obtain from Eqs. (4, 5):

$$\left(\frac{da(t)}{dt}\right)^2 = \frac{1-\Omega_X-\Omega_R}{a(t)} + \frac{\Omega_X}{a(t)^{1+3w_X}} + \frac{\Omega_R}{a(t)^2} - \Omega_G\left(a(t)^2 + \frac{1}{2a(t)^4} - \frac{3}{2a(t)^2}\right), \tag{6}$$

$$\frac{d^2a(t)}{dt^2} = -\frac{1-\Omega_X-\Omega_R}{2a(t)^2} - \frac{(1+3w_X)}{2a(t)^{2+3w_X}} - \frac{\Omega_R}{a(t)^3} - \Omega_G\left(a(t) - \frac{1}{a(t)^5}\right), \tag{7}$$

where $t = H_0(\tau - \tau_0)$, $a(t)$ is the scaling factor normalized to its present value, $\Omega_R = \frac{8\pi\rho_R}{3H_0^2}$ is the density parameter for the radiation (including relativistic neutrinos) with $w_R = 1/3$, $\Omega_X = \frac{8\pi\rho_X}{3H_0^2}$ is the density parameter for the X-matter with unknown negative w_X, $\Omega_G = \frac{m^2}{6H_0^2}$ is the density parameter for the gravitons. For the "usual" matter with density parameter $\Omega_M = \frac{8\pi\rho_M}{3H_0^2}$ we supposed $w_M = 0$. Also, we used the cosmic sum rule $\Omega_M + \Omega_R + \Omega_X = 1$ and took into consideration that the graviton's term doesn't contribute to this rule. As the estimation of w_X can be used the extremal values of the so-called weak energy condition $\rho + p \geq 0$ and strong energy condition $\rho + 3p \geq 0$ [12].

3 Discussion

The dynamical properties of the system (6, 7) can be found by the search of the zeros of the right-hand sides of these equations. Then from the Eq. 6 we have the following polynomial equation for the minimal a_{min} and maximal a_{max} scaling factors of the universe:

$$2(\Omega_X a^{3-3w} - \Omega_G a^6) + 2(1-\Omega_R-\Omega_X)a^3 + (2\Omega_r + 3\Omega_G)a^2 - \Omega_G = 0 \tag{8}$$

The dependences of a_{min} and a_{max} on Ω_G are shown in Fig. 1 for fixed $\Omega_R \approx 9.1 \cdot 10^{-5}$ [12]. As it is known [2, 3, 4], $a_{min} \neq 0$ for nonzero m (curve 1). For $a_{min} < 10^{-4} \approx 1/\Omega_R$ (that is the approximate value of scaling factor at the end of the radiation epoch) the dependence of the minimal scaling factor on Ω_X (and Ω_M) is negligible and we can write for this value:



Fig. 1. The logarithms of the minimal (curve 1) and maximal (curves 2-5) scaling factors in the dependence on the logarithm of the graviton's critical density. Ω_X = 0.65 (1-4), 0 (5), w = -1 (2), -2/3 (3), -1/3 (4)

$$a_{\min} \approx \sqrt{\frac{\Omega_G}{2\Omega_R + 3\Omega_G}}. \tag{9}$$

But the original feature of our model is the strong dependence of a_{max} on the presence and state of X-matter. If its equation of state corresponds to the pure Λ-term (w = -1), there exist the infinitely expanding solutions of Eqs. (6, 7) for $\Omega_G < \Omega_X$ (curve 2, see also the asymptotically growing term in Eq. 8). The transition to more "moderate" w (curves 3, 4) results in the recollapsing behavior of the universe with decreasing a_{max} as result of $|w|$ decrease. The value of a_{max} in the absence of X-matter is shown by curve 5. For fixed w the growth of Ω_X from 0.5 to 0.8 (see Table 1) increases a_{max} but this dependence is reduced by $|w|$ decrease.

Accelerated (decelerated) character of the universe's expansion is defined by the sign of the polynomial, which is the modified right-hand side of Eq. 7:

$$\Omega_X a^{3-3w}(1+3w) + 2\Omega_G a^6 + (1-\Omega_X-\Omega_R)a^3 \\ +2\Omega_R a^2 - 2\Omega_G \tag{10}$$

The roots of this polynomial give the values of the critical scaling factor a_{cr} (Fig. 2) determining the transition from the accelerated (decelerated) to the decelerated (accelerated) expansion.



Fig. 2. The logarithm of the critical scaling factor in the dependence on the logarithm of the graviton's critical density. Ω_X = 0.8 (1), 0.65 (2, 4, 5), 0.5 (3), w = -1 (1 - 3), -2/3 (4), -1/3 (5). Curve 6 shows a_{min}, arrows are directed into regions of the decelerated expansion

It is possible to distinguish following scenarios of the universe's evolution:

I. Deceleration and recollapse. The initially accelerated expansion changes into deceleration with subsequent recollapse. Such behavior takes a place for $\Omega_G > \Omega_X$ in the case of w = -1 (right-hand side group of curves 1 - 3), for Ω_G lying on the right of lower curve 4 in the case of w = -2/3 and for all Ω_G in the case of w = -1/3 (curve 5).

II. Eternal acceleration. For w = -1 there is the region of the simple accelerated expansion. This region increases as result of the Ω_X increase (region between left-hand side and right-hand side groups of curves 1 - 3).

III. Loitering expansion without recollapse. This is a complicated scenario existing for w = -1 and small Ω_G (left-hand side group of curves 1 - 3). The initially accelerated expansion changes into deceleration, which changes into an eternal acceleration not long before the present time. The region of the deceleration increases as result of the Ω_X decrease (transition from the left-hand side curve 1 to 3).

IV. Loitering expansion with recollapse. For w = -2/3 the behavior is the similar to *III*, but the long acceleration changes into deceleration and recollapse (curves 4). The acceleration can be very long for small Ω_G.



Fig. 3. The scenarios I - V. Ω_G = 0.8 (I), 0.5 (II), 10^{-12} (III), 10^{-3} (IV), 0.62 (V); w = -1 (I - III, V), -2/3 (IV). $\Omega_X = 0.65$. Recollapse for scenario IV is located outside of Fig.

V. Marginally loitering. On the right of the region II for w = -1 there is the narrow region near $\Omega_G = \Omega_X$, where the accelerated expansion changes into decelerated one with the subsequent transition to eternal acceleration.

The examples of the described scenarios are shown in Fig. 3.

So, in the presence of X-matter there exist the scenarios with acceleration far from a_{min} (II - V). As the main criterion for the selection of appropriate scenario we consider the demand of the existence of the radiation dominated epoch. The last restricts the value of Ω_G so as $a_{min} < 10^{-4} \approx 1/\Omega_R$. Then from Eq. 9 one can obtain $\Omega_G < 10^{-11.7}$ that gives for the dimensional graviton's mass $m < 10^{-71} g$ if $H_0 = 68 km\ s^{-1}\ Mps^{-1}$. This estimation is essentially lower than in [4]: $m < 4.5 * 10^{-66} g$, because of the last value was obtained on the basis of criterion $a_{max} > 1$ (i. e. the present epoch corresponds to expansion for $\Omega_X = 0$). Obviously, that our criterion is more rigid.

The obtained estimation allows to prefer the scenarios III or IV for the case of the accelerated expansion. The next step is the calculation of the universe's age, that can be made by the integration of Eq. 6:

$$\tau_0 = \frac{1}{H_0} \int_{a_{\min}}^{1} \frac{da}{\sqrt{A}}, \tag{11}$$

Table 2: Universe's age for scenarios III, IV

Ω_X	$\tau_0(w=-1)$, Gyr	$\tau_0(w=-2/3)$, Gyr
0.5	11.6	11.2
0.65	12.9	12.2
0.8	15	13.8

where

$$A = \frac{1-\Omega_X-\Omega_R}{a} + \frac{\Omega_X}{a^{1+3w}} + \frac{\Omega_R}{a^2} - \Omega_g\left(a^2 + \frac{1}{2a^4} - \frac{3}{2a^2}\right).$$

It is convenient to divide the integration on the two part: radiation dominated epoch and matter dominated epoch. For the first interval one can use the approximation, which produced Eq.9. Then the duration of the radiation dominated epoch is $\sim 10^5$ yr with slow dependence on Ω_G for $a_{min} \ll 1/\Omega_R$. The further integration for the matter dominated epoch gives the results, which are presented in Table 2 ($H_0 = 68 km\ s^{-1}\ Mps^{-1}$). The obtained values of the universe's age agree with the modern observational data (see Table 1) and don't depend on the graviton's mass.

4 Conclusion

The introducing of the Λ-term in the Lagrangian of RTG demands $\Lambda = -m^2$ that contributes to the deceleration of the universe's expansion. However, modern observations suggest the accelerated expansion at the present epoch. We showed that without modification of the field equation [8] the insertion of X-matter with the equation of the state, which is close to the extremal weak energy condition $\rho + p = 0$ extends the class of the cosmological models in framework of RTG. As result, there exist the cosmological scenarios, which agree with the observational data. We demonstrated, that in the case of $m < 10^{-71} g$ and $-1 \leq w \leq -2/3$ the minimal size of the universe, its age and acceleration at the present epoch don't contradict with the models considered in the framework of GR. But the existing data don't allow to choose between different scenarios because the similar present behavior results in the essential difference in the future: eternal expansion for $w = -1$ (or $m = 0$), and recollapse for $w > -1$. Moreover, the evolution, as rule, behaves the complicated loitering character with alternation of the acceleration and deceleration. That demands the additional investigations in this direction, especially, the search of RTG consequences for the universe's structure formation.

This work was carried out by the use of the computer algebra system Maple 6. We suppose, that the algorithmization of the RTG basics can stimulate the investigations in this direction. The corresponding commented program can be found and downloaded on [11].

Author is Lise Meitner fellow at the Technical University of Vienna (project M611).

References

[1] A.A. Logunov. "Relativistic theory of gravity", Nova Sc. Publication, 1998.

[2] A.A. Logunov, M.A. Mestvirishvili. Relativistic theory of gravitation, Nauka, Moscow, 1989 (in Russian).

[3] S.S. Gerstein, A.A. Logunov, M.A. Mestvirishvili, *Phys. Atomic Nuclei*, **61**, 1420 (1998).

[4] S.S. Gerstein, A.A. Logunov, M.A. Mestvirishvili, arXiv: hep-th/9711147 (1997).

[5] L. M. Krauss, arXiv: astro-ph/0102305 (2001).

[6] A.G. Riess et al. *Astron. Journ.*, **116**, 1009 (1998).

[7] S. Permutter et al. *Astrophys. Journ.*, **517**, 565 (1999).

[8] V.V. Kiselev, arXiv: hep-ph/0106003 (2001).

[9] S.M. Carroll, *Living reviews in relativity*, **4**(1) (2001).
P. Binétruy, arXiv: hep-ph/0005037 (2000).

[10] V.L. Kalashnikov, arXiv: gr-qc/0103023 (2001).

[11] V.L. Kalashnikov, http://www.geocities.com/optomaplev/programs/rtg1.html

[12] J.D. Cohn, arXiv: astro-ph/9807128 (1998).

[13] D.H. Lyth, arXiv: astro-ph/9312022 (1993).